



UNIVERSITI PUTRA MALAYSIA

**BIOACCUMULATION OF COPPER AND ITS EFFECT ON HUMORAL
IMMUNE RESPONSE IN GOLDFISH CARASSIUS AURATUS
LINNAEUS**

MD. ABUL HOSSAIN

FPSS 1990 2

**BIOACCUMULATION OF COPPER AND ITS EFFECT
ON HUMORAL IMMUNE RESPONSE IN GOLDFISH
CARASSIUS AURATUS LINNAEUS**

By

MD. ABUL HOSSAIN

Thesis Submitted in Fulfillment of the Requirements
for the Degree of Master of Science in the Faculty
of Fisheries and Marine Science
Universiti Pertanian Malaysia

October 1990



TO

MY

PARENTS

ACKNOWLEDGEMENTS

My sincere and deepest appreciation to my supervisor Associate Professor Dr. Mohamed Shariff Mohamed Din for his endless supervision, encouragement and unparalleled friendly cooperation during the entire period of study. I will be ever indebted to him.

I am grateful to my co-supervisors Associate Professor Dr. Law Ah Theem and Lecturer Dr. Patimah Ismail for their continuous guidance and fruitful suggestions during the study.

My special gratitude to Dr. Rohana P. Subasinghe, Dr. Hassan Mohd. Daud and Mr. Liew Hock Chark, Faculty of Fisheries and Marine Science, UPM for their help extended to me during the study.

I would also like to thank Mr. Rosli Aslim and Mr. Perumal Kuppan, Faculty of Fisheries and Marine Science, UPM for their technical assistance.

My sincere acknowledgement goes to Bangladesh Agricultural Research Council (BARC) and International Development Research Centre (IDRC), Canada for financial assistance for the programme.



My boundless gratitude to my mother Mrs. Alimunnesa Amin, father-in-law Mr. Enamul Haque, elder brother Mr. Md. Mokbul Hossain, relatives friends and well wishers for their endless inspirations, encouragement and moral support to complete this programme successfully.

Last but not least, my heartfelt gratitude extended to my beloved wife Mrs. Shahana Hossain (Milu), my only son H. M. Shahneowaz (Lemon) and daughter Mou without whose support, encouragement and tolerance this endeavour would not have been a success.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLESviii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONSxvi
ABSTRACTxvii
ABSTRAK	xx
CHAPTER	
I GENERAL INTRODUCTION	1
Sources of Copper in Aquatic Ecosystem.	1
Effect of Copper on Fish Health	3
Research Objectives	4
II REVIEW OF LITERATURE6
Effect of Copper on Immune Response of Fish.6
Effect of Copper on Physiology of Fish.9
Bioaccumulation of Copper in Fish13
Acute Toxicity of Copper to Fish	20
III DETERMINATION OF MEDIAN LETHAL CONCENTRATION OF COPPER TO GOLDFISH <u>CARASSIUS AURATUS</u> LINNAEUS.	23
Introduction23
Materials and Methods	25



Selection and Maintenance of Test Fish . . .	25
Preparation of Test Solution and Test Fish Exposure	26
Measurement of Copper Concentration	27
Method of Estimation of LC ₅₀ Values of Copper	28
Results	29
Discussion	39
 IV BIOACCUMULATION OF COPPER IN VITAL ORGANS AND MUSCLE OF GOLDFISH <u>CARASSIUS</u> <u>AURATUS</u> LINNAEUS.	 45
Introduction	45
Materials and Methods	47
Collection and Maintenance of Test Fish.	47
Preparation of Copper Solution and Test Fish Exposure.	47
Sampling of Test Fish	48
Measurement of Copper.	49
Solvent Extraction Method.	51
Water Quality	52
Results	53
Discussion	66



V	EFFECT OF COPPER BIOACCUMULATION ON HUMORAL IMMUNE RESPONSE IN GOLDFISH <u>CARASSIUS AURATUS</u> LINNAEUS.	72
	Introduction	72
	Materials and Methods	74
	Maintenance of Test Fish and Their Exposure to Copper	74
	Preparation of Antigen from Sheep Red Blood Cells (SRBC).	74
	Copper Treated Test Fish Immunised with Antigen.	75
	Measurement of Antibody Titre	75
	Results	78
	Discussion	81
VI	CONCLUSION AND RECOMMENDATIONS	87
	BIBLIOGRAPHY	91
	APPENDICES	100
	BIOGRAPHICAL SKETCH	118



LIST OF TABLES

Table	Page
1 Comparison of Distribution of Copper in Tissues of Fish and Shellfish Summarised from Available Research Reports	17
2 Estimated LC ₅₀ Values of Copper to Goldfish Using Simple Linear Probit Regression Model $Y = a + bX$	30
3 Mean Actual Treatment Concentration of Copper (ppm) and Mean of Probit Values of Cumulative Percentage Mortality (in Parenthesis) at Different Exposure Time During 120 h LC ₅₀ Test to Goldfish.. . . .	33
4 Mean Cumulative Copper Percentage Loss During 120 h LC ₅₀ Test to Goldfish.	35
5 Mean 24 Hourly Copper Percentage Loss During 120 h LC ₅₀ Test to Goldfish.	36
6 Mean Residual Copper (ppm) Recorded During 120 h LC ₅₀ Test to Goldfish.	37
7 Physicochemical Water Quality Parameters During 120 h Copper LC ₅₀ Test to Goldfish.	38
8 Actual Exposure Concentration of Copper to Goldfish Recorded During the Experiment	54
9 Mean Bioaccumulation of Copper ($\mu\text{g/g}$ dry weight) in Vital Organs and Muscle of Goldfish After Six Weeks Exposure.	56
10 Mean Bioaccumulation of Copper($\mu\text{g/g}$ dry weight) in Vital Organs and Muscle of Goldfish After Eight Weeks Exposure: Where the Fish were Immunised with SRBC at 6th and 7th Week of the Experiment	57



11	Weekly Mean Percentage Copper Bio-accumulation ($\mu\text{g/g}$ dry weight) in Kidney, Liver, Spleen & Muscle in Pre-immunised and Post-immunised Goldfish	64
12	Range of Physicochemical Water Quality Parameters During Eight Weeks Exposure of Goldfish to Copper.	65
13	Serum Antibody Titres of Goldfish Exposed to Copper for Eight Weeks; Where the Fish were Immunised with SRBC at 6th and 7th Week of the Experiment	79
14	Analysis of Variance for Log Serum Antibody Titres of Goldfish Exposed to Copper for Eight Weeks; Where the Fish were Immunised with SRBC at 6th & 7th Week of the Experiment.	101
15	Regression Analysis for Log Serum Antibody Titres of Goldfish Exposed to Copper for Eight Weeks; Where the Fish were Immunised with SRBC at 6th and 7th Week of the Experiment.. . . .	101
16	Regression Output for 12 h LC_{50} Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality	102
17	Regression Output for 24 h LC_{50} Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality	102
18	Regression Output for 48 h LC_{50} Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality.	103
19	Regression Output for 72 h LC_{50} Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality.	103



20	Regression Output for 96 h LC ₅₀ Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality.	104
21	Regression Output for 120 h LC ₅₀ Value of Copper to Goldfish Calculated Using Probit Values Against Percentage Cumulative Mortality.	104
22	Analysis of Variance for Percentage Cumulative Metal Loss During 120 h LC ₅₀ Test of Copper to Goldfish.. . . .	105
23	Mean Percentage Metal Loss During 120 h LC ₅₀ Test of Copper to Goldfish.	105
24	Regression Analysis of Cumulative Metal Loss Within 24 h During 120 h LC ₅₀ Test of Copper to Goldfish.	106
25	Regression Analysis of Cumulative Metal Loss Within 48 h During 120 h LC ₅₀ Test of Copper to Goldfish	106
26	Regression Analysis of Cumulative Metal Loss Within 72 h During 120 h LC ₅₀ Test of Copper to Goldfish	107
27	Regression Analysis of Cumulative Metal Loss Within 96 h During 120 h LC ₅₀ Test of Copper to Goldfish	107
28	Multiple Range Analysis for Percentage Loss of Copper for Exposure Concentration Group During 120 h LC ₅₀ Test to Goldfish.	108
29	Multiple Range Analysis for Percentage Loss of Copper for Exposure Time Group During 120 h LC ₅₀ Test to Goldfish.	108
30	Two Sample T-test for Actual and Expected Sublethal Exposure Concentration of Copper Throughout the Eight Weeks Experimental Period.	108

31	Regression Analysis for Bioaccumulation in Kidney of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	109
32	Regression Analysis for Bioaccumulation in Liver of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	109
33	Regression Analysis for Bioaccumulation in Spleen of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	110
34	Regression Analysis for Bioaccumulation in Muscle of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	110
35	Regression Analysis for Bioaccumulation in Kidney of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	111
36	Regression Analysis for Bioaccumulation in Liver of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	111
37	Regression Analysis for Bioaccumulation in Spleen of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	112
38	Regression Analysis for Bioaccumulation in Muscle of Goldfish Exposed to 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	112
39	Analysis of Variance for Bioaccumulation of Copper in Vital Organs of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	113



40	Analysis of Variance for Bioaccumulation of Copper in Vital Organs of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	113
41	Analysis of Variance for Weekly Rate of Copper Bioaccumulation in Pre-Immunised and Post-Immunised Goldfish.. . . .	113
42	Means for Bioaccumulation of Copper in Kidney, Liver, Spleen and Muscle of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks	114
43	Means for Bioaccumulation of Copper in Kidney, Liver, Spleen and Muscle of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	114
44	Means for Weekly Rate of Copper Percentage Bioaccumulation in Pre-Immunised and Post-Immunised Goldfish.	115
45	Two Sample T-test for Expected and Actual Exposure Level of Copper Prior to 120 h LC_{50} Test to Goldfish.	115
46	Multiple Range Test for Bioaccumulation of Copper in Different Organs of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.. . . .	116
47	Multiple Range Test for Bioaccumulation of Copper at Different Treatment Group Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Six Weeks.	116
48	Multiple Range Test for Bioaccumulation of Copper in Different Organs of Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper	



	for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.	116
49	Multiple Range Test for Bioaccumulation of Copper at Different Treatment Group Goldfish Treated with 10, 30 and 50 $\mu\text{g/l}$ of Copper for Eight Weeks; Where the Fish were Immunised with SRBC at the End of 6th & 7th Week of the Experiment.. . . .	116
50	Multiple Range Test for Weekly Rate of Copper Bioaccumulation (Data Expressed into Arcsine) for Immunisation Factor Group in Pre-Immunised and Post-Immunised Goldfish.	117
51	Multiple Range Test for Weekly Rate of Copper Bioaccumulation (Data Expressed into Arcsine) for Treatment Factor Group in Pre-Immunised and Post-Immunised Goldfish	117
52	Multiple Range Test for Weekly Rate of Copper Bioaccumulation (Data Expressed into Arcsine) for Organs Factor Group in Pre-Immunised and Post-Immunised Goldfish	117



LIST OF FIGURES

Figure		Page
1	Regression Line (unbroken) of Calculated LC_{50} Values of Copper Against Probit Values of Percentage Cumulative Mortality of Goldfish Exposed for Different Durations: Range of 1st and 2nd Pairs of Broken Lines Represent 99% and 95% Confidence Limits Respectively.	31
2	Standard LC_{50} Curve of Copper to Goldfish: Vertical Bar Represents \pm Standard Error.	32
3	Percentage Cumulative Mortality Curves of Goldfish Treated with Copper: Against, Exposure Time (A) and Treatment (B).	34
4	Mean Actual Sublethal Treatment Concentration of Copper During the Experiment	55
5	The Bioaccumulation in Kidney, Liver, Spleen & Muscle of Goldfish Exposed to 10, 30 & 50 $\mu g/l$ of Copper for Six Weeks (A = Unimmunised) and Eight Weeks (B = Immunised)	58
6	Regression Line (unbroken) of Bioaccumulation in Different Organs of Goldfish Exposed to Copper for Six Weeks: Range of 1st and 2nd Pairs of Broken Lines Represents 99% and 95% Confidence Limits Respectively	59
7	Regression Line (unbroken) of Bioaccumulation in Different Organs of Goldfish Exposed to Copper for Eight Weeks; Where Fish were Immunised with SRBC at 6th & 7th Week of the Experiment:	



	Range of 1st and 2nd Pairs of Broken Lines Represents 99% and 95% Confidence Limits Respectively	60
8	Box and Whisker Plots for Bioaccumulation in Goldfish Exposed for Six Weeks: Treat- ment Group (A) & Organ Group (B); Each Box Represents, Mean Range and Vertical Bar \pm Standard Error.	62
9	Box and Whisker Plots for Bioaccumulation in Goldfish Exposed to Copper for Eight Weeks: Where the Fish were Immunised with SRBC at 6th and 7th Week of the Experiment; Treatment Group (A) & Organ Group (B); Each Box Represents, Mean Range and Vertical Bar \pm Standard Error.	63
10	Box and Whisker Plots for Weekly Rate of Copper Bioaccumulation in Pre-immunised and Post-immunised Goldfish. Each Box Represents, Mean Range and Vertical Bar \pm Standard Error.	65
11	Regression Line (unbroken) of Serum Antibody Titres of Goldfish Exposed to Copper for Eight Weeks; Where Fish were Immunised with SRBC at 6th & 7th Week of the Experiment: Range of 1st & 2nd Pairs of Broken Lines Represents 99% and 95% Confidence Limits Respectively.	80



LIST OF ABBREVIATIONS

APDC	= Ammonium Pyrrolidine Dithiocarbamate
APHA	= American Public Health Association
AWWA	= American Water Works Association
DDTC	= Diethyl Ammonium Diethyldithiocarbamate
DO	= Dissolved Oxygen
EIFAC	= European Inland Fisheries Advisory Commission
EPA	= Environmental Protection Agency of U.S.A
FAO	= Food and Agriculture Organisation of U.N
h	= Hours
huRBC	= Human Red Blood Cell
IPNV	= Infectious Pancreatic Necrosis Virus
MATC	= Maximum Allowable Toxicant Concentration
MIBK	= Methyl Isobutyl Ketone
mon.	= Months
PBS	= Phosphate Buffer Saline
S.D	= Sample Standard Deviation
S.E	= Sample Standard Error
SRBC	= Sheep Red Blood Cells
Pre-IM	= Pre-Immunised
Post-IM	= Post-Immunised
V/V	= Volume by Volume
WPCF	= Water Pollution Control Federation of U.S.A
Wt.	= Weight



Abstract of thesis submitted to the Senate of Universiti
Pertanian Malaysia in fulfillment of the requirements
for the degree of Master of Science

**BIOACCUMULATION OF COPPER AND ITS EFFECT
ON HUMORAL IMMUNE RESPONSE IN GOLDFISH
CARASSIUS AURATUS LINNAEUS**

By

MD. ABUL HOSSAIN

December, 1990

Supervisor : Dr. Mohd. Shariff Mohd. Din.
Associate Professor

Faculty : Fisheries and Marine Science

This study reports on the acute toxicity levels, bioaccumulation of copper and their effects on the humoral immune response of goldfish (Carassius auratus). This study also provides information on the levels of bioaccumulation of copper in organs such as kidney, liver and spleen, which plays a role on the humoral antibody production of goldfish under tropical conditions.

The acute toxicity levels of copper to goldfish (10.5 ± 1.43 g) for 12, 24, 48, 72, 96 and 120 hours were estimated as 0.44 ± 0.06 , 0.36 ± 0.04 , 0.20 ± 0.03 ,



0.19 ± 0.02 , 0.15 ± 0.02 and 0.14 ± 0.02 ppm respectively under fresh water static conditions.

Goldfish (17 ± 0.2 g) were exposed to sublethal levels (10, 30 and 50 $\mu\text{g/l}$) of copper for a period of six and eight weeks to facilitate bioaccumulation. The group which was exposed to copper for eight weeks was injected with SRBC at the end of sixth and seventh weeks. This was done to activate the antibody production in test fish and to monitor the effect of copper on humoral antibody response. The bioaccumulation of copper in kidney, liver, spleen and muscle of goldfish was directly proportional to the exposure concentration. The highest bioaccumulation was observed in liver and was followed by spleen, kidney and muscle. Bioaccumulation of copper at 14.19–23.51 $\mu\text{g/g}$ in kidney, 60.28–102.99 $\mu\text{g/g}$ in liver, 15.09–26.30 $\mu\text{g/g}$ in spleen and 2.66–4.11 $\mu\text{g/g}$ in muscle (dry weight) was evident following six weeks exposure. Bioaccumulation of copper in organs of test fish exposed for eight weeks was as follows: kidney 20.39–28.84 $\mu\text{g/g}$, liver 82.01–152.77 $\mu\text{g/g}$, spleen 22.38–39.31 $\mu\text{g/g}$ and muscle 3.39–5.29 $\mu\text{g/g}$ (dry weight).



At the end of eighth week, serum anti-SRBC agglutinating antibody titres in goldfish, which were exposed to copper for eight weeks and subjected to immunisation at sixth and seventh weeks, were partially suppressed when compared to unexposed fish. The suppression of antibody titre was inversely proportional to the exposure concentration of copper. The fish unexposed and exposed to copper at 50 $\mu\text{g/l}$ showed highest 1:256 (mode) and lowest 1:32 (mode) antibody titres respectively. The recorded pre-immunisation heterophilic (nonspecific) antibody titre was 1:4 (mode). Antibody titre of fish unexposed and exposed to copper at different sublethal levels were significantly different.

The weekly rate of bioaccumulation of copper in post-immunised fish was significantly higher when compared to the pre-immunised fish.



Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia, sebagai memenuhi keperluan untuk mendapat Ijazah Master Sains.

BIOAKUMULASI KUPRUM DAN KESANNYA TERHADAP RANGSANGAN IMUN HUMORAL DALAM IKAN MAS CARASSIUS AURATUS LINNAEUS

Oleh

MD. ABUL HOSSAIN

Disember, 1990

Penyelia : Prof. Madya Dr. Mohd. Shariff Mohd. Din.

Fakulti : Perikanan dan Sains Samudera

Kajian ini melaporkan tahap toksisiti akut, bioakumulasi kuprum dan kesan terhadap pengeluaran antibodi humoral dalam ikan mas (C. auratus). Kajian ini juga menyatakan tahap bioakumulasi kuprum dalam organ seperti ginjal, hati dan limpa yang mana memainkan peranan dalam pengeluaran antibodi humoral ikan mas di dalam keadaan tropika.

Tahap toksisiti akut kuprum pada ikan mas (10.5 ± 1.43 g) untuk 12, 24, 48, 72, 96 dan 120 jam masing-masing dianggarkan pada 0.44 ± 0.06 , 0.36 ± 0.04 , 0.20 ± 0.03 , 0.19 ± 0.02 , 0.15 ± 0.02 dan 0.14 ± 0.02 bpj, di dalam keadaan air tawar stasis.

Ikan mas (17 ± 0.2 g) telah didedah kepada tahap subletal (10, 30 dan 50 $\mu\text{g/l}$) kuprum selama enam dan lapan minggu untuk memudahkan bioakumulasi. Kumpulan yang didedahkan kepada kuprum selama lapan minggu telah disuntik dengan SRBC pada akhir minggu ke enam dan ke tujuh. Ini adalah untuk mengaktifkan pengeluaran antibodi dalam ikan yang diuji dan untuk meneliti kesan kuprum terhadap gerak-balas antibodi humoral. Bioakumulasi kuprum dalam ginjal, hati dan limpa adalah berkadaran langsung dengan kepekatan pendedahan. Bioakumulasi tertinggi telah dilihat dalam hati, dituruti oleh limpa, ginjal dan otot. Bioakumulasi kuprum dalam ginjal ikan mas adalah 14.19-23.51 $\mu\text{g/g}$, hati 60.28-102.99 $\mu\text{g/g}$, limpa 15.09-26.30 $\mu\text{g/g}$ dan otot 2.66-4.11 $\mu\text{g/g}$ berat kering apabila didedahkan selama enam minggu. Bioakumulasi kuprum dalam ikan mas yang didedahkan selama lapan minggu adalah seperti berikut: ginjal 20.39-28.84 $\mu\text{g/g}$, hati 82.01-152.77 $\mu\text{g/g}$, limpa 22.38-39.31 $\mu\text{g/g}$ dan otot 3.39-5.29 $\mu\text{g/g}$ (berat kering).

Pada minggu ke lapan, titer serum anti-SRBC antibodi pengaglutinatan dalam ikan mas yang didedahkan pada kuprum selama lapan minggu dan diimunkan dalam minggu ke enam dan ke tujuh, didapati separa tertekan jika dibandingkan dengan ikan yang tidak terdedah. Penekanan titer antibodi adalah berkadar songsang dengan kepekatan pendedahan kuprum.

Ikan yang tidak didedah dan yang didedah kepada kuprum (50 µg/l) masing-masing menunjukkan titer antibodi tertinggi, 1:256 (mod) dan terendah 1:32 (mod). Titer antibodi heterofilik (bukan khusus) sebelum imunisasi dirakamkan pada 1:4 (mod). Titer antibodi ikan yang tidak didedah dan didedahkan kepada kuprum pada paras subletal yang berlainan, adalah berbeza keertian, di antara satu sama lain.

Kadar mingguan bioakumulasi kuprum dalam ikan pasca-imun adalah lebih tinggi keertian berbanding dengan ikan praimun.

CHAPTER I

GENERAL INTRODUCTION

Sources of Copper in Aquatic Ecosystem

Copper is one of the most commonly available, toxic and bioaccumulable heavy metals present in water. With the rapid industrialisation of the present world, and the indiscriminate discharge of copper oriented products/complexans, the metal is continuously increasing in both fresh and sea water ecosystems. The industrial sources of copper from widely used major industrial products are fertilizers, algicides, pesticides, therapeutic agents, bronze, wiring, storage batteries, electrical goods, etc.

Copper is also widely distributed in nature as rocks and earth crust. Copper is introduced into the aquatic ecosystem as a result of weathering of soils and rocks from volcanic eruption and also from a variety of human activities, including mining process. Contamination of copper from the industries occur when the metals



or other substances that contain copper contaminants are discharged into the aquatic ecosystem (Mance, 1987; Nriagu, 1979; EIFAC/FAO, 1976). The sources of copper rich water is usually from contaminated mine water or run-off from aged spoil heaps (Mance, 1987). Thus the ecosystem is being constantly polluted by unextracted copper compounds from the soil and from the abandoned industrial products containing copper (Nriagu, 1979).

Copper can exist in the form of at least 5000 complexans in water (EIFAC/FAO, 1976). Copper when present in combination with other heavy metals especially zinc, becomes more harmful to aquatic organisms. Copper is safe for normal biochemical life processes if its presence is below 5 $\mu\text{g/l}$ in water. The atmospheric loading of copper into most terrestrial and aquatic ecosystem now exceeds the combined copper requirement of flora and fauna for their normal biochemical life processes (Kelly et al., 1975). In Kelang Estuary, Malaysia, the mean total and dissolved copper content was 10 $\mu\text{g/l}$ and 4.3 $\mu\text{g/l}$ respectively (Law and Singh, 1986).